KYUNGPOOK Math. J. 62(2022), 323-331
https://doi.org/10.5666/KMJ.2022.62.2.323
pISSN 1225-6951 eISSN 0454-8124
© Kyungpook Mathematical Journal

# Pseudohermitian Curvatures on Bounded Strictly Pseudoconvex Domains in $\mathbb{C}^2$

Aeryeong Seo
*Department of Mathematics, Kyungpook National University, Daegu 41566, Republic of Korea*
*e-mail* : aeryeong.seo@knu.ac.kr

Abstract. In this paper, we present a formula for pseudohermitian curvatures on bounded strictly pseudoconvex domains in $\mathbb{C}^2$ with respect to the coefficients of adapted frames given by Graham and Lee in [3] and their structure equations. As an application, we will show that the pseudohermitian curvatures on strictly plurisubharmonic exhaustions of Thullen domains diverges when the points converge to a weakly pseudoconvex boundary point of the domain.

## 1. Introduction

Let $(M, \theta)$ be a pseudohermitian manifold of real dimension $2n+1$, i.e. $M$ is an integrable, nondegenerate, real hypersurface in $\mathbb{C}^{n+1}$ with CR structure $HM \subset TM$ so that $\theta$ is a 1-form satisfying $\theta(X) = 0$ for any $X \in HM$ and $d\theta$ can be expressed by

$$d\theta = ih_{\alpha\bar{\beta}}\theta^{\alpha} \wedge \theta^{\bar{\beta}}$$

where $(h_{\alpha\bar{\beta}})$ is a positive definite $n \times n$ matrix with respect to a local basis $\theta^1, \ldots, \theta^n$ for $HM$. The equivalence problem for such pseudohermitian structures was studied first by Chern using Cartan's method and later it turned out that it was related to the pseudoconformal invariants analyzed by Chern-Moser [1] and Tanaka [4]. In [5], Webster showed that there exists a natural connection in the bundle $H^{1,0}M$ adapted to a pseudohermitian structure, where $H^{1,0}M$ denotes the eigenspace of the endomorphism $J\colon HM \otimes \mathbb{C} \to HM \otimes \mathbb{C}$ satisfying $J \circ J = -I$, which defines the CR structure of $M$, with an eigenvalue $\sqrt{-1}$. Moreover he gave an expression of pseudoconformal curvature tensor in terms of pseudohermitian curvature tensor.

In this paper, we present an expression of pseudohermitian curvatures on bounded strictly pseudoconvex domains in $\mathbb{C}^2$ with respect to the coefficients of adapted frames given by Graham-Lee in [3] and their structure equations. As an

Received February 25, 2022; revised June 16, 2022; accepted June 17, 2022.
2020 Mathematics Subject Classification: 32V05, 53B15.
Key words and phrases:Pseudohermitian manifolds, Thullen domains.
This research was supported by Kyungpook National University Research Fund, 2019.

application, we will show that the pseudohermitian curvatures on strictly plurisubharmonic exhaustions of Thullen domains diverge when the point converges to the weakly pseudoconvex boundary point of the domain. Our main result is

**Main Theorem.** *For a Thullen domain*

$$\Omega_m := \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^{2m} + |z_2|^2 < 1\}$$

*with $m > 1$, the pseudohermitian curvatures of the exhaustions*

$$\Omega_{m,\epsilon} := \{(z_1, z_2) \in \mathbb{C}^2 : \phi(z_1, z_2) = \epsilon\}$$

*of $\Omega_m$ with $\phi = -K(z,z)^{-m/(2m+1)}$ diverge as the point tends to weakly pseudoconvex boundary points along the complex line $\{z_1 = 0\}$ in $\Omega_m$.*

## 2. Pseudohermitian Curvatures on Strictly Pseudoconvex Domains

In this section, we calculate the pseudohermitian curvature explicitly in terms of the coefficient of an adapted coframe for a strictly pseudoconvex bounded domain $\Omega := \{(z_1, z_2) \in \mathbb{C}^2 : \phi(z_1, z_2) < 0\}$ where $\phi$ is a defining function of $\Omega$. Let $\{\theta^0, \theta^1\}$ be the adapted coframe and $\{X^0, X^1\}$ its dual frame given by Graham-Lee ([3]) satisfying the following.

$$\begin{aligned}
d\theta^0 &= -\theta^1 \wedge \overline{\theta^1} - r\theta^0 \wedge \overline{\theta^0} \\
d\theta^1 &= \theta^1 \wedge \omega - iA\theta^0 \wedge \overline{\theta^1} + \frac{r}{2}\theta^0 \wedge \theta^1 + \frac{r}{2}\overline{\theta^0} \wedge \theta^1 - X^1 r\theta^0 \wedge \overline{\theta^0} \\
d\omega &= R\theta^1 \wedge \theta^{\bar{1}} + \left( i\left(X^{\bar{1}}A - 2\omega(X^{\bar{1}}\bar{A})\right) - \frac{3}{2}(X^1 r)\right)\theta^1 \wedge \theta^{\bar{0}} \\
&\quad - \left( i\left(X^1\bar{A} - 2\omega(X^1 A)\right) + \frac{3}{2}(X^{\bar{1}} r)\right)\theta^0 \wedge \theta^{\bar{1}} + \frac{3}{2}(X^1 r)\theta^0 \wedge \theta^1 \\
&\quad - \frac{3}{2}(X^{\bar{1}} r)\theta^{\bar{0}} \wedge \theta^{\bar{1}} + \left(\frac{1}{2}\Delta_b r - |A|^2\right)\theta^0 \wedge \theta^{\bar{0}}
\end{aligned} \tag{2.1}$$

Here $R$ denotes the pseudohermitian curvature and $\Delta_b := -r_\alpha{}^\alpha - r^\alpha{}_\alpha$ denotes the sublaplacian of the pseudohermitian structure. In coordinates of $\mathbb{C}^2$, let

$$\theta^0 := A_1 dz_1 + A_2 dz_2, \quad \theta^1 := B_1 dz_1 + B_2 dz_2,$$

$$\omega := C_1 dz_1 + C_2 dz_2 - \overline{C_1} d\overline{z}_1 - \overline{C_2} d\overline{z}_2, \tag{2.2}$$

and

$$X^0 = \frac{1}{D}\left(B_2\frac{\partial}{\partial z_1} - B_1\frac{\partial}{\partial z_2}\right), \quad X^1 = \frac{1}{D}\left(-A_2\frac{\partial}{\partial z_1} + A_1\frac{\partial}{\partial z_2}\right)$$

where $D := A_1B_2 - A_2B_1$ and $A_j := \frac{\partial \phi}{\partial z_j}$ for each $j = 1, 2$.

**Proposition 2.1.** *Let $\Omega = \{(z_1, z_2) \in \mathbb{C}^2 : \phi(z_1, z_2) < 0\}$ be a strictly pseudoconvex domain with a defining function $\phi$. Then the pseudohermitian curvature is given by*

$$R = \frac{1}{|B_1|^2}\left(2Re\left(\left(i(X^1\bar{A} - 2\omega(X^{\bar{1}}\bar{A})) + \frac{3}{2}(X^{\bar{1}}r)\right)A_1\overline{B_1} - \frac{\partial C_1}{\partial \overline{z}_1}\right) - \left(\frac{1}{2}\Delta_b r - |A|^2\right)|A_1|^2\right).$$

*Proof.* By (2.1), we have

$$\begin{aligned} d\theta^0 &= -\theta^1 \wedge \overline{\theta^1} - r\theta^0 \wedge \overline{\theta^0} \\ &= -(B_1dz_1 + B_2dz_2) \wedge (\overline{B_1dz_1 + B_2dz_2}) \\ &\quad - r(A_1dz_1 + A_2dz_2) \wedge (\overline{A_1dz_1 + A_2dz_2}) \\ &= -(|B_1|^2 + r|A_1|^2)dz_1 \wedge d\overline{z}_1 - (B_1\overline{B}_2 + rA_1\overline{A}_2)dz_1 \wedge d\overline{z}_2 \\ &\quad - (B_2\overline{B}_1 + rA_2\overline{A}_1)dz_2 \wedge d\overline{z}_1 - (|B_2|^2 + r|A_2|^2)dz_2 \wedge d\overline{z}_2. \end{aligned} \tag{2.3}$$

On the other hand, since

$$\begin{aligned} d\theta^0 &= d(A_1dz_1 + A_2dz_2) \\ &= -\frac{\partial A_1}{\partial \overline{z}_1}dz_1 \wedge d\overline{z}_1 - \frac{\partial A_1}{\partial \overline{z}_2}dz_1 \wedge d\overline{z}_2 - \frac{\partial A_2}{\partial \overline{z}_1}dz_2 \wedge d\overline{z}_1 - \frac{\partial A_2}{\partial \overline{z}_2}dz_2 \wedge d\overline{z}_2, \end{aligned} \tag{2.4}$$

by comparing equations (2.3) and (2.4) we have

$$|B_1|^2 + r|A_1|^2 = \frac{\partial A_1}{\partial \overline{z}_1}, \tag{2.5}$$

$$B_1\overline{B}_2 + rA_1\overline{A}_2 = \frac{\partial A_1}{\partial \overline{z}_2}, \tag{2.6}$$

and

$$|B_2|^2 + r|A_2|^2 = \frac{\partial A_2}{\partial \overline{z}_2}. \tag{2.7}$$

Since we have

$$\begin{aligned} |B_1|^2|B_2|^2 &= \left(\frac{\partial A_1}{\partial \overline{z}_1} - r|A_1|^2\right)\left(\frac{\partial A_2}{\partial \overline{z}_2} - r|A_2|^2\right) \\ &= \left(\frac{\partial A_1}{\partial \overline{z}_2} - rA_1\overline{A}_2\right)\left(\frac{\partial \overline{A_1}}{\partial z_2} - rA_2\overline{A}_1\right) \end{aligned}$$

by (2.5), (2.7), (2.6), we obtain

$$r = \frac{\left|\frac{\partial A_1}{\partial \overline{z}_2}\right|^2 - \frac{\partial A_1}{\partial \overline{z}_1}\frac{\partial A_2}{\partial \overline{z}_2}}{A_1\overline{A}_2\frac{\partial \overline{A_1}}{\partial z_2} + A_2\overline{A}_1\frac{\partial A_1}{\partial \overline{z}_2} - |A_2|^2\frac{\partial A_1}{\partial \overline{z}_1} - |A_1|^2\frac{\partial A_2}{\partial \overline{z}_2}}. \tag{2.8}$$

By substituting (2.8) into (2.5) and (2.7), we have

$$|B_1|^2 = \frac{A_1\overline{A_2}\frac{\partial A_1}{\partial \overline{z}_1}\frac{\partial \overline{A_1}}{\partial z_2} + A_2\overline{A_1}\frac{\partial A_1}{\partial \overline{z}_1}\frac{\partial A_1}{\partial \overline{z}_2} - |A_2|^2\left(\frac{\partial A_1}{\partial \overline{z}_1}\right)^2 - |A_1|^2\left|\frac{\partial A_1}{\partial \overline{z}_2}\right|^2}{A_1\overline{A}_2\frac{\partial \overline{A_1}}{\partial z_2} + A_2\overline{A}_1\frac{\partial A_1}{\partial \overline{z}_2} - |A_2|^2\frac{\partial A_1}{\partial \overline{z}_1} - |A_1|^2\frac{\partial A_2}{\partial \overline{z}_2}}, \tag{2.9}$$

and

$$|B_2|^2 = \frac{A_1\overline{A}_2\frac{\partial \overline{A_1}}{\partial z_2}\frac{\partial A_2}{\partial \overline{z}_2} + A_2\overline{A}_1\frac{\partial A_1}{\partial \overline{z}_2}\frac{\partial A_2}{\partial \overline{z}_2} - |A_1|^2\left(\frac{\partial A_2}{\partial \overline{z}_2}\right)^2 - |A_2|^2\left|\frac{\partial A_1}{\partial \overline{z}_2}\right|^2}{A_1\overline{A}_2\frac{\partial \overline{A_1}}{\partial z_2} + A_2\overline{A}_1\frac{\partial A_1}{\partial \overline{z}_2} - |A_2|^2\frac{\partial A_1}{\partial \overline{z}_1} - |A_1|^2\frac{\partial A_2}{\partial \overline{z}_2}}. \tag{2.10}$$

By (2.1) and a straightforward calculation we obtain

$$\begin{aligned}
d\theta^1 &= \theta^1\wedge\omega - iA\theta^0\wedge\overline{\theta^1} + \frac{r}{2}\theta^0\wedge\theta^1 + \frac{r}{2}\overline{\theta^0}\wedge\theta^1 - X^1r\theta^0\wedge\overline{\theta^0}\\
&= (B_1dz_1 + B_2dz_2)\wedge(C_1dz_1 + C_2dz_2 - \overline{C_1}d\overline{z}_1 - \overline{C_2}d\overline{z}_2)\\
&\quad - iA(A_1dz_1 + A_2dz_2)\wedge\overline{(B_1dz_1 + B_2dz_2)}\\
&\quad + \frac{r}{2}(A_1dz_1 + A_2dz_2)\wedge(B_1dz_1 + B_2dz_2)\\
&\quad + \frac{r}{2}\overline{(A_1dz_1 + A_2dz_2)}\wedge(B_1dz_1 + B_2dz_2)\\
&\quad - X^1r(A_1dz_1 + A_2dz_2)\wedge\overline{(A_1dz_1 + A_2dz_2)}\\
&= \left(B_1C_2 - B_2C_1 + \frac{r}{2}A_1B_2 - \frac{r}{2}A_2B_1\right)dz_1\wedge dz_2\\
&\quad + \left(-B_1\overline{C_1} - iAA_1\overline{B_1} - \frac{r}{2}\overline{A_1}B_1 - (X^1r)|A_1|^2\right)dz_1\wedge d\overline{z}_1\\
&\quad + \left(-B_1\overline{C_2} - iAA_1\overline{B_2} - \frac{r}{2}\overline{A_2}B_1 - (X^1r)A_1\overline{A_2}\right)dz_1\wedge d\overline{z}_2\\
&\quad + \left(-B_2\overline{C_1} - iAA_2\overline{B_1} - \frac{r}{2}\overline{A_1}B_2 - (X^1r)A_2\overline{A_1}\right)dz_2\wedge d\overline{z}_1\\
&\quad + \left(-B_2\overline{C_2} - iAA_2\overline{B_2} - \frac{r}{2}\overline{A_2}B_2 - (X^1r)|A_2|^2\right)dz_2\wedge d\overline{z}_2.
\end{aligned} \tag{2.11}$$

On the other hand, since

$$\begin{aligned}
d\theta^1 &= d(B_1dz_1 + B_2dz_2)\\
&= -\frac{\partial B_1}{\partial \overline{z}_1}dz_1\wedge d\overline{z}_1 - \frac{\partial B_1}{\partial z_2}dz_1\wedge dz_2 - \frac{\partial B_1}{\partial \overline{z}_2}dz_1\wedge d\overline{z}_2\\
&\quad - \frac{\partial B_2}{\partial z_1}dz_2vdz_1 - \frac{\partial B_2}{\partial \overline{z}_1}dz_2\wedge d\overline{z}_1 - \frac{\partial B_2}{\partial \overline{z}_2}dz_2\wedge d\overline{z}_2,
\end{aligned} \tag{2.12}$$

by comparing equations (2.11) and (2.12), we obtain

$$B_1C_2 - B_2C_1 + \frac{r}{2}A_1B_2 - \frac{r}{2}A_2B_1 = -\frac{\partial B_1}{\partial z_2} + \frac{\partial B_2}{\partial z_1}, \tag{2.13}$$

$$B_1\overline{C_1} + iAA_1\overline{B_1} + \frac{r}{2}\overline{A_1}B_1 + (X^1r)|A_1|^2 = \frac{\partial B_1}{\partial \overline{z}_1}, \tag{2.14}$$

$$B_1\overline{C_2} + iAA_1\overline{B_2} + \frac{r}{2}\overline{A_2}B_1 + (X^1r)A_1\overline{A_2} = \frac{\partial B_1}{\partial \overline{z}_2},$$

$$B_2\overline{C_1} + iAA_2\overline{B_1} + \frac{r}{2}\overline{A_1}B_2 + (X^1r)A_2\overline{A_1} = \frac{\partial B_2}{\partial \overline{z}_1},$$

and

$$B_2\overline{C_2} + iAA_2\overline{B_2} + \frac{r}{2}\overline{A_2}B_2 + (X^1r)|A_2|^2 = \frac{\partial B_2}{\partial \overline{z}_2}. \tag{2.15}$$

By considering $B_2 \times (2.14) - B_1 \times (2.15)$, we have

$$A = \frac{1}{i\overline{B_1}D}\left(\frac{\partial B_1}{\partial \overline{z}_1}B_2 - \frac{\partial B_2}{\partial \overline{z}_1}B_1 - (X^1r)\overline{A_1}D\right). \tag{2.16}$$

By substituting (2.16) into (2.14), we obtain

$$\overline{C_1} = \frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_1} - \frac{1}{D}\frac{A_1}{B_1}\left(\frac{\partial B_1}{\partial \overline{z}_1}B_2 - \frac{\partial B_2}{\partial \overline{z}_1}B_1\right) - \frac{r}{2}\overline{A_1}.$$

Similarly by considering $B_2 \times (2.15) - B_1 \times (2.15)$ we have

$$A = \frac{1}{i\overline{B_2}D}\left(\frac{\partial B_1}{\partial \overline{z}_2}B_2 - \frac{\partial B_2}{\partial \overline{z}_2}B_1 - (X^1r)\overline{A_2}D\right) \tag{2.17}$$

and by substituting (2.17) into (2.15), we have

$$\overline{C_2} = \frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_2} - \frac{1}{D}\frac{A_1}{B_1}\left(\frac{\partial B_1}{\partial \overline{z}_2}B_2 - \frac{\partial B_2}{\partial \overline{z}_2}B_1\right) - \frac{r}{2}\overline{A_2}.$$

Now using the expression (2.1) and (2.2), we obtain

$$\begin{aligned}
d\omega &= R(B_1dz_1 + B_2dz_2) \wedge \overline{(B_1dz_1 + B_2dz_2)} \\
&\quad + \left(i\left(X^{\bar{1}}A - 2\omega(X^{\bar{1}}\bar{A})\right) - \frac{3}{2}(X^1r)\right)(B_1dz_1 + B_2dz_2) \wedge \overline{(A_1dz_1 + A_2dz_2)} \\
&\quad - \left(i\left(X^1\bar{A} - 2\omega(X^1A)\right) + \frac{3}{2}(X^{\bar{1}}r)\right)(A_1dz_1 + A_2dz_2) \wedge \overline{(B_1dz_1 + B_2dz_2)} \\
&\quad + \frac{3}{2}(X^1r)(A_1dz_1 + A_2dz_2) \wedge (B_1dz_1 + B_2dz_2) \\
&\quad - \frac{3}{2}(X^{\bar{1}}r)\overline{(A_1dz_1 + A_2dz_2)} \wedge \overline{(B_1dz_1 + B_2dz_2)} \\
&\quad + (\Delta_b r - |A|^2)(A_1dz_1 + A_2dz_2) \wedge \overline{(A_1dz_1 + A_2dz_2)} \\
&= d(C_1dz_1 + C_2dz_2 - \overline{C_1}d\overline{z}_1 - \overline{C_2}d\overline{z}_2).
\end{aligned} \tag{2.18}$$

By comparing the coefficient of $dz_1 \wedge d\overline{z}_1$ in the equation (2.18), one has

$$-\frac{\partial C_1}{\partial \overline{z}_1} - \frac{\partial \overline{C_1}}{\partial z_1} = R|B_1|^2 + \left( i\left( X^{\bar{1}}A - 2\omega(X^{\bar{1}}\bar{A}) \right) - \frac{3}{2}(X^1 r) \right) B_1 \overline{A_1} \\ - \left( i\left( X^1 \bar{A} - 2\omega(X^1 A) \right) + \frac{3}{2}(X^{\bar{1}} r) \right) A_1 \overline{B_1} + \left( \frac{1}{2}\Delta_b r - |A|^2 \right) |A_1|^2.$$

As a result, we complete the proof. □

## 3. Pseudohermitian Curvature of Thullen Domains

For $m \in \mathbb{N}$, let

$$\Omega_m := \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^{2m} + |z_2|^2 < 1\}$$

be the Thullen domain in $\mathbb{C}^2$ with $m \geq 1$. The diagonal of the Bergman kernel of $\Omega_m$ is given by

$$K(z,z) = \frac{1}{m\pi^2} \frac{(m+1)\left(1-|z_2|^2\right)^{1/m} - (1-m)|z_1|^2}{(1-|z_2|^2)^{2-1/m}\left((1-|z_2|^2)^{1/m} - |z_1|^2\right)^3}.$$

For the detail, see [2]. Then $\phi(z, \bar{z}) := -K(z,z)^{-m/(2m+1)}$ gives a defining function of $\Omega_m$.

For a point $a \in \Delta := \{z \in \mathbb{C} : |z| < 1\}$, denote by $\psi_a$ an automorphism of $\Omega_m$ obtained by

$$\psi_a : (z_1, z_2) \mapsto \left( \frac{(1-|a|^2)^{1/2m} z_1}{(1-\overline{a}z_2)^{1/m}}, \frac{z_2 - a}{1 - \overline{a}z_2} \right).$$

Note that $\psi_{-a}$ is the inverse of $\psi_a$. Then we have

$$\mathrm{Aut}(\Omega_m) = \left\{ (z_1, z_2) \mapsto \left( e^{i\theta_1} \frac{(1-|a|^2)^{\frac{1}{2m}} z_1}{(1-\overline{a}z_2)^{\frac{1}{m}}}, e^{i\theta_2} \frac{z_2 - a}{1 - \overline{a}z_2} \right) : \theta_1, \theta_2 \in \mathbb{R}, a \in \Delta \right\}$$

and $W := \{(0, w) : |w| = 1\} \subset \partial\Omega_m$ is the set of weakly pseudoconvex boundary points of $\Omega_m$ for $m > 1$. Note that $\partial\Omega_m \setminus W$ is the set of strictly pseudoconvex boundary points.

From now on, we will find the pseudohermitian curvature of the exhaustion

$$\Omega_{m,\epsilon} := \{(z_1, z_2) \in \mathbb{C}^2 : \phi(z_1, z_2) = \epsilon\}$$

with $\epsilon \ll 1$. In particular, we will observe the behavior of the pseudohermitian curvature when $(0, z_2) \to (0, w)$ with $|w| = 1$, i.e. weakly pseudoconvex points of

$\Omega_m$. At $z=(0,z_2)$, we have the following:

$$
\begin{aligned}
&K(z,z)=\frac{1}{\pi^2}\frac{m+1}{m}\frac{1}{(1-|z_2|^2)^{2+\frac{1}{m}}},\\
&\phi=-\left(\frac{1}{\pi^2}\frac{m+1}{m}\right)^{\frac{-m}{2m+1}}(1-|z_2|^2),\\
&A_1=\frac{\partial A_2}{\partial z_1}=\frac{\partial A_1}{\partial z_2}=\frac{\partial A_2}{\partial \overline{z}_1}=\frac{\partial A_1}{\partial \overline{z}_2}=\frac{\partial A_1}{\partial z_1}=\frac{\partial^2 A_1}{(\partial \overline{z}_1)^2}=0\\
&A_2=-\phi\frac{\overline{z}_2}{1-|z_2|^2}=\left(\frac{1}{\pi^2}\frac{m+1}{m}\right)^{\frac{-m}{2m+1}}\overline{z}_2,\quad \frac{\partial A_2}{\partial \overline{z}_2}=\left(\frac{1}{\pi^2}\frac{m+1}{m}\right)^{\frac{-m}{2m+1}}.
\end{aligned}
$$

Hence at $z=(0,z_2)$, by straightforward calculation we have

$$
\begin{aligned}
&r=\frac{1}{|A_2|^2}\frac{\partial A_2}{\partial \overline{z}_2}=\left(\frac{1}{\pi^2}\frac{m+1}{m}\right)^{\frac{m}{2m+1}}\frac{1}{|z_2|^2},\\
&|B_1|^2=\frac{\partial A_1}{\partial \overline{z}_1}=\frac{\partial^2\phi}{\partial z_1\partial\overline{z}_1}=-\frac{2m}{m+1}\frac{1}{(1-|z_2|^2)^{1/m}}\phi,\quad B_2=0,\\
&A=\frac{1}{iA_2\overline{B_1}}\frac{\partial B_1}{\partial \overline{z}_1},\quad \overline{C_1}=\frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_1},\quad \overline{C_2}=\frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_2}-\frac{r}{2}\overline{A_2},\\
&R=\frac{-2}{|B_1|^2}\mathrm{Re}\left(\frac{\partial C_1}{\partial \overline{z}_1}\right).
\end{aligned}
\tag{3.1}
$$

By differentiating the equation (2.5) with $\frac{\partial}{\partial\overline{z}_1}$, we obtain

$$
\begin{aligned}
\frac{\partial B_1}{\partial\overline{z}_1}\overline{B_1}+B_1\frac{\partial\overline{B_1}}{\partial\overline{z}_1}&=\frac{\partial}{\partial\overline{z}_1}\left(\frac{\partial A_1}{\partial\overline{z}_1}-r|A_1|^2\right)\\
&=\frac{\partial^2 A_1}{\partial^2\overline{z}_1}-\frac{\partial r}{\partial\overline{z}_1}|A_1|^2-r\frac{\partial|A_1|^2}{\partial\overline{z}_1}.
\end{aligned}
\tag{3.2}
$$

Since $A_1=0$ and $\frac{\partial A_1}{\partial z_1}=0$ at $z=(0,z_2)$, we have

$$
\frac{\partial B_1}{\partial\overline{z}_1}\overline{B_1}+B_1\frac{\partial\overline{B_1}}{\partial\overline{z}_1}=0,
$$

and hence

$$
\frac{1}{B_1}\frac{\partial B_1}{\partial\overline{z}_1}=-\frac{1}{\overline{B_1}}\frac{\partial\overline{B_1}}{\partial\overline{z}_1}.
$$

By differentiating the equation (3.2) with $\frac{\partial}{\partial z_1}$, we obtain

$$
\begin{aligned}
&\frac{\partial^2 B_1}{\partial z_1\partial\overline{z}_1}\overline{B_1}+\left|\frac{\partial B_1}{\partial\overline{z}_1}\right|^2+\left|\frac{\partial B_1}{\partial z_1}\right|^2+B_1\frac{\partial^2\overline{B_1}}{\partial z_1\partial\overline{z}_1}\\
&=\frac{\partial^3 A_1}{\partial z_1\partial^2\overline{z}_1}-\frac{\partial^2 r}{\partial z_1\partial\overline{z}_1}|A_1|^2-\frac{\partial r}{\partial\overline{z}_1}\frac{\partial|A_1|^2}{\partial z_1}-\frac{\partial r}{\partial z_1}\frac{\partial|A_1|^2}{\partial\overline{z}_1}-r\frac{\partial^2|A_1|^2}{\partial z_1\partial\overline{z}_1}.
\end{aligned}
$$

Evaluating the above equation at $z = (0, z_2)$ gives

$$2\mathrm{Re}\left(\frac{1}{B_1}\frac{\partial^2 B_1}{\partial z_1 \partial \overline{z}_1}\right) + 2\left|\frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_1}\right|^2 = \frac{1}{|B_1|^2}\left(\frac{\partial^3 A_1}{\partial z_1 \partial^2 \overline{z}_1} - r\left|\frac{\partial A_1}{\partial \overline{z}_1}\right|^2\right). \tag{3.3}$$

Since

$$\begin{aligned}\frac{\partial \overline{C_1}}{\partial z_1} &= \frac{\partial}{\partial z_1}\left(\frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_1} - \frac{1}{D}\frac{A_1}{B_1}\left(\frac{\partial B_1}{\partial \overline{z}_1}B_2 - \frac{\partial B_2}{\partial \overline{z}_1}B_1\right) - \frac{r}{2}\overline{A_1}\right)\\ &= -\frac{1}{B_1^2}\frac{\partial B_1}{\partial z_1}\frac{\partial B_1}{\partial \overline{z}_1} + \frac{1}{B_1}\frac{\partial^2 B_1}{\partial z_1 \partial \overline{z}_1} - \frac{r}{2}\frac{\partial \overline{A_1}}{\partial z_1}\\ &= \left|\frac{1}{B_1}\frac{\partial B_1}{\partial z_1}\right|^2 + \frac{1}{B_1}\frac{\partial^2 B_1}{\partial z_1 \partial \overline{z}_1} - \frac{r}{2}\frac{\partial \overline{A_1}}{\partial z_1},\end{aligned}$$

by Proposition 2.1 and equations (3.3), (3.1) we obtain

$$\begin{aligned}R &= \frac{-2}{|B_1|^2}\left(\left|\frac{1}{B_1}\frac{\partial B_1}{\partial \overline{z}_1}\right|^2 + \mathrm{Re}\left(\frac{1}{B_1}\frac{\partial^2 B_1}{\partial z_1 \partial \overline{z}_1}\right) - \frac{r}{2}\frac{\partial \overline{A_1}}{\partial z_1}\right)\\ &= \frac{-1}{|B_1|^2}\left(\frac{1}{|B_1|^2}\left(\frac{\partial^3 A_1}{\partial z_1 \partial^2 \overline{z}_1} - r\left|\frac{\partial A_1}{\partial \overline{z}_1}\right|^2\right) - r\frac{\partial \overline{A_1}}{\partial z_1}\right)\\ &= \frac{-1}{|B_1|^4}\frac{\partial^3 A_1}{\partial z_1 \partial^2 \overline{z}_1} + 2r\frac{1}{|B_1|^2}\frac{\partial A_1}{\partial \overline{z}_1}\\ &= \frac{-1}{|B_1|^4}\frac{\partial^4 \phi}{\partial^2 z_1 \partial^2 \overline{z}_1} + 2r,\end{aligned} \tag{3.4}$$

$$\begin{aligned}\frac{\partial \phi}{\partial z_1} &= \phi\frac{\partial}{\partial z_1}\log(-\phi) = -p\phi\frac{\partial}{\partial z_1}\log K\\ &= -p\phi\frac{\partial}{\partial z_1}\log\left((m+1)\left(1-|z_2|^2\right)^{1/m} - (1-m)|z_1|^2\right)\\ &\quad + 3p\phi\frac{\partial}{\partial z_1}\log\left(\left(1-|z_2|^2\right)^{1/m} - |z_1|^2\right)\\ &= -p\phi\overline{z}_1\left(\frac{-(1-m)}{(m+1)(1-|z_2|^2)^{\frac{1}{m}} - (1-m)|z_1|^2} + \frac{3}{(1-|z_2|^2)^{\frac{1}{m}} - |z_1|^2}\right)\end{aligned}$$

with $p = \frac{m}{2m+1}$, and

$$\begin{aligned}\frac{\partial^2 \phi}{\partial^2 z_1} &= p^2\overline{z}_1^2\phi\left(\frac{-(1-m)}{(m+1)(1-|z_2|^2)^{\frac{1}{m}} - (1-m)|z_1|^2} + \frac{3}{(1-|z_2|^2)^{\frac{1}{m}} - |z_1|^2}\right)^2\\ &\quad - p\phi\overline{z}_1^2\left(\frac{-(1-m)^2}{\left((m+1)(1-|z_2|^2)^{\frac{1}{m}} - (1-m)|z_1|^2\right)^2} + \frac{3}{\left((1-|z_2|^2)^{\frac{1}{m}} - |z_1|^2\right)^2}\right).\end{aligned}$$

Let

$$M := \frac{-(1-m)}{(m+1)(1-|z_2|^2)^{\frac{1}{m}} - (1-m)|z_1|^2} + \frac{3}{(1-|z_2|^2)^{\frac{1}{m}} - |z_1|^2}$$

and

$$N := \frac{-(1-m)^2}{\Big((m+1)(1-|z_2|^2)^{\frac{1}{m}} - (1-m)|z_1|^2\Big)^2} + \frac{3}{\Big((1-|z_2|^2)^{\frac{1}{m}} - |z_1|^2\Big)^2}.$$

Then

$$\frac{\partial^2 \phi}{\partial^2 z_1} = p^2 \overline{z}_1^2 M^2 \phi - p\phi \overline{z}_1^2 N.$$

Hence at $z = (0, z_2)$, we have

$$\frac{\partial^4 \phi}{\partial^2 z_1 \partial^2 \overline{z}_1} = 2p\phi(pM^2 - N)$$

and by (3.4)

$$R = -\frac{(3m+1)(m-1)}{m(2m+1)} \frac{1}{\phi} + 2r. \tag{3.5}$$

As a result we obtain our main theorem.

# References


[1] S. S. Chern and J. K. Moser, *Real hypersurfaces in complex manifolds*, Acta Math., **133**(1974), 219–271.

[2] J. P. D'Angelo, *A note on the Bergman kernel*, Duke Math. J., **45(2)**(1978), 259–265.

[3] C. R. Graham and J. M. Lee, *Smooth solutions of degenerate Laplacians on strictly pseudoconvex domains*, Duke Math. J., **57(3)**(1988), 697–720.

[4] N. Tanaka, *On the pseudo-conformal geometry of hypersurfaces of the space of n complex variables*, J. Math. Soc. Japan, **14**(1962), 397–429.

[5] S. M. Webster, *Pseudo-Hermitian structures on a real hypersurface*, J. Differential Geometry, **13(1)**(1978), 25–41.